LAKE: POVERTY P (BIG) (VLMP 31 )

TOWN: NEWFIELD COUNTY: YORK

MIDAS: 157
TRUE BASIN: 1
SAMPLE STATION:

1

TRUE BASIN CHARACTERISTICS

SURFACE AREA: 60.0 ha. (148.3 a.)

FLUSHING RATE: 2.72 flushes/yr.

VOLUME: 1625266.3 cu. m. (1318 ac.-ft.)

DIRECT DRAINAGE AREA: 4.76 sq. km. (1.84 sq. mi.)

WHOLE LAKE INFORMATION

MAX. DEPTH: 5 m. (17 ft.)

MEAN DEPTH: 3 m. (10 ft.)

DELORME ATLAS #: 02

USGS QUAD: LIMERICK


IFW REGION A: Sebago Lake (Gray)


IIW RESIGN A. Sebago Lake (Siay)

IFW FISH. MANAGMENT: Warmwater & Coldwater

PLEASE NOTE THE FOLLOWING: The SAMPLE STATION # refers to the location sampled. The term TRUE BASIN is used to define areas within a lake that are separated by shallow reefs or shoals and therefore function as separate lakes. There are approximately 50 lakes in the state that have more than 1 True Basin. True Basin Characteristics are now being included in the first section of these reports to enable users of the Phosphorous Loading Methodology to better evaluate the data. If there is no data for a particular True Basin, True Basin Characteristics must be obtained from the DEP. POVERTY P (BIG) has 1 True Basin(s).

## SECCHI DISK TRANSPARENCY GRAPHS:





Note: 2010 graphs may indicate multiple readings taken on a given day.

## SUMMARY OF CHEMICAL AND TROPHIC STATE PARAMETERS:

[\* indicates that Secchi disk was visable at bottom of lake (or one reading used in calculation was visable)].

	MEAN	MEAN	MEAN	MEAN															
	COLOR	pН	ALK	COND.	TOTAL	PHOS.	MEANS	(ppb)	SECCH	I DISK	(m.)		CHLORO	PHYLL	A(ppb)	TROP	HIC ST	ATE IN	DICES
	(SPU)		(mg/l)	(uS	EPI	SURF	BOT.	PRO.								EPI	PHOS		
YEAR				/cm)	CORE	GRAB	GRAB	GRAB	MIN.	MEAN	MAX.	<u>N</u> _	MIN.	MEAN	MAX.	<u>C</u>	G	SEC	CHL
1978	- 20	6.80	7.0	30	9	-	-	N 175	4.1	4.1	4.1	1	2.2	2.2	2.2	-	-	-	-
1979	25	-		-	-	-	-	-	-	-	-		-	-	-	-	-	-	
1997	14	6.91	7.0	42	10	-	-	-	3.9*	4.2*	4.4*	1	2.5	2.5	2.5	-	-	-	-
2000	15	-	9.0	42	10	-	-	-	4.4*	4.4*	4.4*	1	4.1	4.1	4.1	-	-	-	-
2007	- 1		-	-	-	1-1	-	-	4.2*	4.6*	4.7	5	-	=	-	-	-	-	-
2008	24	6.95	5.6	42	-	6	-	-	4.3*	4.7*	4.8*	4	-	-	-	-	-	-	-
2009	-	a a 740			-	-	-	-	4.5*	4.7*	4.8*	5	-	-	-	-	-	-	
2010	-	-	-	_	-	-	-	-	4.5*	4.6*	4.9*	4	-	-	-	-	-	-	-
SUMMARY:	20	6.88	7.2	39	10	6	-	+:	3.9	4.5*	4.9*	7	2.2	2.9	4.1	-	-	-	-

LAKE: POVERTY P (BIG) (VLMP 31 )

MIDAS: 157
\*TRUE BASIN: 1

TOWN: NEWFIELD COUNTY: YORK

\*SAMPLE STATION: 1

LATE SUMMER TEMPERATURE / DISSOLVED OXYGEN PROFILES:

		SAMPLE DATE	1
DEPTH	08/21/78	08/21/97	08/25/00
m	°C ppm	°C ppn	°C ppm
0.0	27.6 99.9	23.1 8.1	21.5 9.0
1.0	1- 1-	23.1 8.0	21.0 9.0
2.0	_(0., 5.1)	23.1 7.9	21.0 9.0
3.0	- " -	23.1 7.9	21.0 8.9
MARK TA	5 91	22 1 7 0	21 0 8 9

ramida rii - W. -

avealth set also as a bladering an

k Indix W in it

## WATER QUALITY SUMMARY

## POVERTY POND (BIG), NEWFIELD

MIDAS: 0157, Sample Station # 1

The Maine Department of Environmental Protection (ME-DEP) and the Volunteer Lake Monitoring Program (VLMP) have collaborated in the collection of lake data to evaluate water quality, track algal blooms, and determine water quality trends. This dataset does not include bacteria, mercury, or nutrients other than phosphorus.

Water quality monitoring data for Poverty Pond has been collected since 1978. During this period, 4 years of basic chemical information was collected in addition to Secchi Disk Transparencies (SDT). In summary, the water quality of Poverty Pond is considered average, based on measures of SDT, total phosphorus (TP), and Chlorophyll-a (Chla). The potential for nuisance algal blooms on Poverty Pond is low to moderate.

Water Quality Measures: Poverty Pond is a non-colored lake (average color 29 SPU) with an average SDT of 4.5\*m (14.3\*ft). (The asterisk\* indicates that the SDT readings sometimes were visible on the bottom of the pond. If the pond were deeper, the SDT readings would be greater.) The range of water column TP for Poverty Pond is 9 to 10 parts per billion (ppb) with an average of 10 ppb. Chla ranges from 2.2 to 4.1 ppb with an average of 2.9 ppb. Recent dissolved oxygen (DO) profiles show little DO depletion in deep areas of the lake. The potential for phosphorus to leave the bottom sediments and become available to algae in the water column (internal loading) is low. Oxygen levels below 5 parts per million stress certain cold water fish and a persistent loss of oxygen may eliminate or reduce habitat for sensitive cold water species.

See ME-DEP Explanation of Lake Water Quality Monitoring Report for measured variable explanations. Additional lake information can be found on the Internet at <a href="http://www.lakesofmaine.org/">http://www.lakesofmaine.org/</a> and/or <a href="http://www.maine.gov/dep/blwq/lake.htm">http://www.maine.gov/dep/blwq/lake.htm</a>, or telephone the ME-DEP at 207-287-3901 or the VLMP at 207-783-7733.

Filename: pove0157, Revised: 03/03, By: jhm Updated: 2/11, By: jp